


# Dynapenic abdominal obesity is associated with negative clinical outcomes in older patients with type 2 diabetes: a prospective cohort study

Mileni Vanti Beretta<sup>a,b</sup> , Ticiana da Costa Rodrigues<sup>a,b</sup> , Thais Steemburgo<sup>a,b</sup> 

<sup>a</sup>Universidade Federal do Rio Grande do Sul – Porto Alegre (RS), Brazil.

<sup>b</sup>Hospital de Clínicas de Porto Alegre – Porto Alegre (RS), Brazil.

## Correspondence data

Thais Steemburgo – Rua Ramiro Barcelos, 2400, 2º andar – CEP: 90035-003 – Porto Alegre (RS), Brazil.  
E-mail: tsteemburgo@gmail.com

**Received on:** Jul 24, 2024

**Editor decision on:** Aug 27, 2024

**Accepted on:** Oct 12, 2024

**Handling Editor:** Ana Patricia Navarrete-Reyes

**How to cite this article:** Beretta MV, Rodrigues TC, Steemburgo T. Dynapenic abdominal obesity is associated with negative clinical outcomes in older patients with type 2 diabetes: a prospective cohort study. *Geriatr Gerontol Aging.* 2024;18:e0000230. [https://doi.org/10.53886/gga.e0000230\\_EN](https://doi.org/10.53886/gga.e0000230_EN)

**Copyright:** © 2024 Beretta et al. This open-access article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



## Abstract

**Objective:** To evaluate the association between dynapenic abdominal obesity (DAO) and the following negative outcomes in older patients with type 2 diabetes (T2D): prolonged length of stay (LOS), readmission within 30 days, in-hospital mortality, and mortality within 1 year after hospital discharge.

**Methods:** This prospective cohort study included hospitalized older patients with T2D. DAO was defined as a combination of abdominal obesity (high waist circumference [WC]:  $\geq 102$  cm for men and  $\geq 88$  cm for women]) and dynapenia (reduced hand grip strength [HGS]:  $< \leq 27$  kg for men and  $< 16$  kg for women]). The association between clinical outcomes and DAO was evaluated using multivariate analyses adjusted for confounders. The Kaplan-Meier curve was used to compare 1-year survival in the presence of DAO.

**Results:** We included 309 patients with T2D (mean age  $73.3 \pm 6.4$  years; 50.5% female; 32.4% with DAO). In multivariate analyses, patients with DAO had a 5.29- and 4.71-fold increase in LOS ( $\geq 14$  days) and 1-year mortality than those without DAO, respectively. Moreover, patients with DAO had a higher risk of 1-year mortality (log-rank test,  $p < 0.05$ ).

**Conclusions:** Older patients with T2D and DAO are more likely to have prolonged hospitalization and 1-year mortality compared to those without DAO.

**Keywords:** abdominal obesity; older adults; type 2 diabetes.

## INTRODUCTION

Abdominal obesity (AO) and dynapenia are two distinct health conditions related to aging and lifestyle.<sup>1</sup> AO can contribute to decreased muscle strength by the action of inflammatory and endocrine mechanisms that occur due to muscle fat infiltration.<sup>1</sup> Dynapenia is characterized by decreased muscle strength and functionality, mostly in older adults.<sup>2</sup> Individuals with dynapenia have functional limitations that further hinder their performance of basic activities, worsening their quality of life.<sup>3</sup> The combination of dynapenia and AO can better predict functional disability than either condition by itself.<sup>4</sup> Therefore, this recently recognized condition was named dynapenic abdominal obesity (DAO).<sup>5</sup>

DAO is emerging as a major risk factor, as recent studies have shown it to be associated with increased risk of numerous negative outcomes, including fractures<sup>4,5</sup>, cognitive impairment,<sup>6</sup> worsening disability, and mortality.<sup>7,8</sup> Moreover, DAO is particularly concerning in patients with type 2 diabetes (T2D), as it can exacerbate the adverse effects associated with diabetes.<sup>1</sup> Isolated OA is a common feature of T2D, and is linked to increased insulin resistance and a higher risk of cardiovascular diseases.<sup>1</sup> Moreover, older patients with T2D often experience a decline in muscle strength owing to poor glycemic control, chronic inflammation, and other metabolic disturbances.<sup>9,10</sup> This reduction in muscle strength – dynapenia – can impair mobility and increase the risk of falls.<sup>11</sup> Thus, DAO can negatively impact the health of older adults with T2D, increasing systemic inflammation and insulin resistance and causing greater difficulty in controlling blood glucose levels.<sup>6-8</sup>

Previous studies in different populations have indicated a relationship between low muscle strength, prolonged hospital stay, and mortality.<sup>12-14</sup> Likewise, in older adults without diabetes, DAO was associated with a higher risk of hospitalization – but not with mortality.<sup>15</sup>

Despite the importance of DAO as a significant health problem, specifically in older people with T2D due to its connection with various adverse clinical outcomes, studies in this group of individuals remain scarce. Thus, this study aimed to assess the association of DAO with the following negative clinical outcomes in hospitalized older patients with T2D: prolonged length of stay (LOS), readmission within 30 days, in-hospital mortality, and mortality within 1 year after hospital discharge.

## METHODS

This was the second part of a cohort study that included older individuals with and without T2D.<sup>16</sup> The study protocol was

approved by the institutional Research Ethics Committee (opinion #150068) and all participants provided written informed consent, as per the recommendations established by the Declaration of Helsinki.

Participants aged  $\geq 60$  years with T2D, who had been admitted from July 2015 to December 2017, were considered eligible. Individuals in the intensive care unit, in palliative care, with neurological sequelae, bedbound, and those using enteral nutrition who were unable to communicate or complete the anthropometric measurements were excluded. Figure 1 shows a flow diagram of the patient selection process.

Data were collected within 48 hours of hospital admission at the patient's bedside by a trained researcher using a standardized protocol.

General features, including age and sex, were collected from electronic records. Ethnicity was self-reported by the patient or next of kin upon hospital admission and recorded in participants' electronic medical records. Self-reported smoking and alcohol consumption were prompted by the yes/now questions "Are you a smoker or former smoker?" and "Do you usually consume alcoholic beverages?" Self-reported physical activity was prompted by the question: "Do you practice any physical activity? (yes/no) If so, what activity, how many times a week, and for how long?" Reason for hospitalization and clinical data (such as duration of diabetes, presence of comorbidities, and use of antihyperglycemic agents) were collected from medical records. Disease severity was scored according to the age-adjusted Charlson comorbidity index (CCI).<sup>17</sup> Laboratory measurements (HbA1C, albumin, and C-reactive protein levels) were also collected from medical records.

The participants were weighed and measured within 48 hours of admission by a trained registered dietitian. Body mass index (BMI) was estimated as individuals' weight (kg) divided by their height (cm) squared. BMI was classified according to the WHO criteria<sup>18</sup> as underweight (BMI  $< 18.5$  kg/m<sup>2</sup>), normal weight (BMI  $\geq 18.5$ – $24.99$  kg/m<sup>2</sup>), or overweight (BMI  $\geq 25$  kg/m<sup>2</sup>). Malnutrition was evaluated by the Mini Nutritional Assessment Long Form (MNA-LF).<sup>19</sup> The MNA-LF is a specific tool for older adults that consists of 18 components grouped into four aspects: anthropometric data, general status, dietary habits, self-perceived health, and nutritional state.<sup>19</sup> The cutoff points of the MNA-LF classified participants as well-nourished (24–30 points), at risk of malnutrition (17–23.5 points), or malnourished ( $< 17$  points).<sup>19</sup>

Abdominal obesity (AO) was assessed by the waist circumference (WC), measured and recorded in centimeters using a 143 cm long inelastic tape (1 mm resolution), with




FIGURE 1. Flow diagram of patient selection.

the patient standing upright, at the approximate midpoint between the lower margin of the last rib and the upper bound of the iliac crest. The cut-off points used to classify high WC were  $\geq 88$  cm for women and  $\geq 102$  cm for men, as per WHO criteria.<sup>18</sup>

Dynapenia was assessed by handgrip strength (HGS), measured using a hydraulic hand dynamometer (Saehan<sup>®</sup>) to record the maximum strength patients' non-dominant hand could exert for three seconds. For HGS measurement, patients were instructed to remain seated with their elbows flexed at  $90^\circ$  along the sides of the body. The highest measurement was used for analysis. Values  $< 16$  kg for women and  $< 27$  kg for men were considered as reduced muscle strength, indicating dynapenia.<sup>20</sup>

Dynapenic abdominal obesity (DAO) was defined by the co-occurrence of abdominal obesity (WC  $\geq 102$  cm for men and  $\geq 88$  cm for women)<sup>18</sup> and dynapenia (HGS  $< 27$  kg for men and  $< 16$  kg for women).<sup>20</sup>

Cognitive status was assessed in all participants using the Mini-Mental State Examination (MMSE), which yields a score of 0 to 30 points. The cut-off point was set according

to educational attainment:  $< 25$  points for those with 1 to 4 years of education; 26.5 for those with 5 to 8 years of education; 28 for those with 9 to 11 years of education; and 29 for those with  $\geq 11$  years of education.<sup>21</sup>

The length of stay (LOS) was estimated in days from the date of admission to the date of discharge. Prolonged hospitalization ( $\geq 14$  days) was considered according to patients' median LOS. Hospital readmission and mortality data were collected by telephone contact every 3 months with patients or close family members or friends until 1 year after discharge. When telephone contact was not possible, the patient's electronic medical record was consulted. The main causes of mortality were collected from medical records.

This is a secondary analysis of a previously published prospective clinical study of older adults with and without diabetes who had been admitted to a university hospital.<sup>16</sup>

The Kolmogorov–Smirnov test was used to assess the normality of distribution of quantitative variables. Continuous variables are expressed as means  $\pm$  standard deviations or medians (interquartile ranges) as appropriate;

categorical variables are expressed as absolute (n) and relative (%) frequencies.

For the analysis, participants were divided into four groups by presence or absence of dynapenia (D) and abdominal obesity (AO): Group 1, D (-) /AO (-); Group 2, AO (+); Group 3, D (+); and Group 4, DAO (+). Kruskal–Wallis and ANOVA tests were used as necessary.

Multivariate analyses with 95% confidence intervals (CI) were performed considering the following negative clinical outcomes: LOS ≥ 14 days (categorized by a median of 14 days considering the data distribution in the sample), 1-year hospital readmission (logistic regression), in-hospital mortality, and 1-year mortality (Cox regression). Moreover, independent models were constructed in the multivariate analysis considering negative clinical outcomes as independent predictors. Stepwise regression was used to select the most important covariates based on their contribution to model fit. All models were adjusted for the Charlson comorbidity index (adjusted for age), surgical procedures during hospital stay, time since the diagnosis of diabetes, and the presence of malnutrition.

The Kaplan–Meier curve (log-rank  $p < 0.05$ ) was used to assess the 1-year survival of patients with DAO. All statistical analyses were performed using SPSS, version 25.0 (SPSS Inc., Chicago, IL, USA). Statistical significance was set at  $p < 0.05$ .

## RESULTS

Table 1 describes the general characteristics of hospitalized older patients with T2D. This study included 309 participants with a mean age of  $73.3 \pm 6.4$  years, of whom 50.5% ( $n = 156$ ) were women and 72% ( $n = 223$ ) were white. Regarding lifestyle factors, about 14% of participants smoked and consumed alcohol, and 84.8% reported performing no physical activities. The main reason for hospitalization was surgical intervention (29.8%;  $n = 92$ ). Among clinical and laboratory characteristics, 74.6% of patients had hypertension, with a median duration of diabetes of 10 (2–15) years, median HbA1c of 7% (6.4–8.5%), and 35.5% ( $n = 110$ ) were on antihyperglycemic agents. Analysis showed a high CCI adjusted for median age (three points, range one to four), normal albumin values (3.7 mg/dl, range 3.2 to 4.1), and high CRP values (26 mg/dl, range 7.10–93).

On nutritional evaluation at hospital admission, mean BMI was  $27.9 \pm 5.3$  kg/m<sup>2</sup>; 68.2% of individuals were overweight (BMI ≥ 25 kg/m<sup>2</sup>) and 17.8% ( $n = 55$ ) had malnutrition according to the MNA-LF. In the assessment of abdominal obesity, 78% of patients had high WC, with a

**TABLE 1.** Characteristics of 309 hospitalized older adults with type 2 diabetes.

Demographic characteristics (%)	
Age (years)	73.3 ± 6.4
Sex (female)	156 (50.5)
Ethnicity/skin color (white)	223 (72)
Behavioral characteristics (%)	
Smoking (yes)	45 (14.6)
Alcohol consumption (yes)	43 (14)
Sedentary lifestyle (yes)	262 (84.8)
Reason for hospitalization (%)	
Surgical intervention	92 (29.8)
Cancer	62 (20)
Liver and bile duct infections	87 (28.1)
Falls	46 (14.8)
Diabetes complications	14 (4.5)
Lung diseases	8 (2.6)
Clinical and laboratory characteristics	
Comorbidities (%)	
Hypertension	232 (74.6)
Cardiovascular disease	42 (19.6)
Duration of diabetes (years)	10 (2–15)
HbA1C (%)	7 (6.4–8.5)
Antihyperglycemic agents	110 (35.5)
CCI	3 (1–4)
Albumin (mg/dL)	3.70 (3.2–4.1)
C-reactive protein (mg/dL)	26 (7.10–93)
Nutritional parameters (%)	
BMI (kg/m <sup>2</sup> )	27.9 ± 5.3
≥ 25 kg/m <sup>2</sup>	211 (68.2)
Malnutrition (MNA-LF)	55 (17.8)
Abdominal obesity	
WC, women (cm)	96.6 ± 13.7
WC, men (cm)	98.8 ± 12.5
High WC*	241 (78)
Dynapenia (%)	
HGS, women (kg)	15.6 ± 5.6
HGS, men (kg)	23.7 ± 7.1
Low HGS <sup>†</sup>	153 (49.5)
Dynapenic abdominal obesity <sup>‡</sup>	100 (32.4)
Cognition (%)	
Cognitive impairment (MMSE)	78 (25.2)
Negative outcomes (%)	
LOS (days)	14 (8–20)
LOS ≥ 14 days	156 (50.5)
One-year hospital readmission	123 (40)
In-hospital mortality	19 (6.1)
One-year mortality	67 (21.7)
Causes of mortality (%)	
Cancer	31 (10)
Sepsis and shock	25 (8.1)
Cardiovascular diseases	21 (6.8)
Generalized complications	9 (3)

HbA1C: Glycated Hemoglobin Test; CCI: Charlson comorbidity index; BMI: body mass index; MNA-LF: mini nutritional assessment long form; WC: waist circumference; HGS: handgrip strength; MMSE: Mini Mental Status Exam; LOS: length of hospital stay.

Data expressed as mean ± standard deviation, median (interquartile range), or n (%).

\*High WC: men, ≥102 cm; women, ≥ 88 cm<sup>18</sup>; <sup>†</sup>Low HGS: men, < 27 kg; women, < 16 kg<sup>20</sup>; <sup>‡</sup>Dynapenic abdominal obesity is defined as the combination of high WC and low HGS.

mean of  $96.6 \pm 13.7$  cm in women and  $98.8 \pm 12.5$  cm in men. Mean HGS was  $15.6 \pm 5.6$  kg for women and  $23.7 \pm 7.1$  kg for men. Low HGS, denoting dynapenia, occurred in 49.5% ( $n = 153$ ) of individuals. Overall, 38.2% of patients were diagnosed with dynapenic abdominal obesity.

Evaluation of cognition by MMSE showed that 25.2% of participants had cognitive impairment. Of the negative outcomes, median LOS was 14 (8 – 20) days, 50.5% of participants remained hospitalized for  $\geq 14$  days, overall, in-hospital mortality was 6.1%, and 1-year mortality was 21.7%. The main causes of mortality (in-hospital and 1 year after discharge) were cancer (10%), sepsis and shock (8.1%), cardiovascular diseases (6.8%), and generalized complications (3%).

Table 2 shows patients' characteristics stratified by the presence of dynapenia and DAO. Of the 309 older individuals with T2D, 9.38% ( $n = 29$ ) had no AO or dynapenia but 27.5% ( $n = 85$ ) had AO; 30.7% ( $n = 95$ ) had dynapenia; and

32.4% ( $n = 100$ ) had DAO. AO was more prevalent in women than in men (62%;  $p = 0.001$ ). Older people with dynapenia had had diabetes for longer than those in other groups ( $p = 0.004$ ). Regarding nutritional variables, about 45% of individuals with AO had a  $\text{BMI} \geq 25 \text{ kg/m}^2$ . Moreover, 24.5% ( $n = 23$ ) of individuals with dynapenia had malnutrition according to the MNA-LF.

Table 3 shows the associations of dynapenia and AO with negative clinical outcomes, adjusted for CCI, surgical procedures during hospital stay, time since diagnosis of diabetes, and malnutrition. Older individuals with dynapenia alone and those with DAO had higher odds of LOS  $\geq 14$  days (odds ratio [OR] = 1.61 [95%CI 1.12 – 3.87] and 2.50 [95%CI 1.05 – 5.95], respectively). Furthermore, older adults with T2D and DAO had a 4.51-fold increased risk of 1-year mortality than older adults with T2D without DAO. Kaplan-Meier survival analysis confirmed this finding (Figure 2).

**TABLE 2.** Characteristics of 309 older adults hospitalized with type 2 diabetes, stratified by presence of dynapenia and abdominal obesity status.


	D (-)/AO (-) (n = 29; 9.38%)	AO (+) (n = 85; 27.5%)	D (+) (n = 95; 30.7%)	DAO (+) (n = 100; 32.4%)
Demographic variables				
Age (years)	74.9 $\pm$ 4.4	75.8 $\pm$ 4.8	75.6 $\pm$ 4.6	75.7 $\pm$ 4.6
Sex (female)*	10 (34.4)	59 (69.4)	25 (26.3)	62 (62)
Ethnicity/skin color (white)	20 (69)	64 (75.3)	68 (71.6)	71 (71)
Behavioral variables				
Smoking (yes)	5 (15.2)	15 (17.6)	13 (13.7)	12 (12)
Alcohol consumption (yes)	6 (20.7)	9 (10.6)	14 (14.7)	14 (14)
Sedentary lifestyle (yes)	24 (82.8)	76 (89.4)	80 (84.2)	82 (82)
Clinical conditions				
Surgical procedures (yes)	11 (37.9)	31 (36.5)	30 (31.6)	20 (20)
Hypertension (yes)	22 (66.7)	97 (80.2)	23 (65.7)	89 (75.4)
Duration of diabetes (years) <sup>†</sup>	1.5 (1 – 10)	5 (1 – 15)	15 (5 – 15)	10 (5 – 12)
HbA1C (%)	7.15 $\pm$ 1.83	7.44 $\pm$ 1.99	7.59 $\pm$ 2.24	7.67 $\pm$ 2.11
Albumin (mg/dL)	3.73 $\pm$ 0.75	3.72 $\pm$ 0.66	3.47 $\pm$ 0.54	3.73 $\pm$ 0.73
C-reactive protein (mg/dL)	33 (1.4 – 185)	20 (0 – 82.5)	24.4 (1.35 – 59.7)	44.5 (1.90 – 124)
Nutritional evaluation				
BMI ( $\geq 25 \text{ kg/m}^2$ ) <sup>‡</sup>	12 (41.4)	38 (44.7)	23 (24.6)	36 (36)
Malnutrition (MNA-LF)	6 (20)	11 (13)	23 (24.5)	15 (15)
Cognition				
Cognitive impairment (MMSE)	5 (17.2)	18 (21.2)	25 (26.3)	30 (30)
Negative outcomes				
LOS $\geq 14$ days	10 (34.5)	39 (45.9)	54 (56.8)	56 (53)
One-year hospital readmission	12 (41.4)	39 (45.9)	38 (40)	34 (34)
One-year mortality <sup>§</sup>	3 (10.3)	10 (11.2)	23 (24.2)	31 (31)

D: Dynapenia; AO: Abdominal obesity; DAO: Dynapenic abdominal obesity; HbA1C: Glycated Hemoglobin Test; BMI: Body mass index; MNA-LF: Mini nutritional assessment long form; MMSE: Mini Mental Status Exam; LOS: Length of hospital stay. Data expressed as mean  $\pm$  standard deviation, median (interquartile range), or n (%). One-way analysis of variance (ANOVA) was used for continuous variables and the Kruskal-Wallis test for categorical variables. \* $p = 0.001$ ; <sup>†</sup> $p = 0.004$ ; <sup>‡</sup> $p < 0.001$ ; <sup>§</sup> $p = 0.006$ .

**TABLE 3.** Association of dynapenia and abdominal obesity with negative clinical outcomes: multivariate analysis.

Dependent variables	Prolonged LOS (≥ 14 days)		One-year hospital readmission		In-hospital mortality		One-year mortality	
Independent variables	OR* (95%CI)	p-value	OR* (95%CI)	p-value	HR† (95%CI)	p-value	HR† (95%CI)	p-value
D (-)/AO (-) (reference)	1	-	-	-	-	-	-	-
AO (+)	1.79 (0.72 – 4.42)	0.205	1.38 (0.59 – 3.22)	0.44	0.96 (0.33 – 2.80)	0.94	2.87 (0.84 – 9.77)	0.90
D (+)	1.61 (1.12 – 3.87)	0.029	0.86 (0.29 – 2.53)	0.79	1.94 (0.55 – 6.77)	0.29	1.34 (0.36 – 4.99)	0.65
DAO (+)	2.50 (1.05 – 5.95)	0.038	0.98 (0.42 – 2.30)	0.97	1.75 (0.62 – 4.98)	0.46	4.51 (1.35 – 15.36)	0.014

All models of multivariate analysis were adjusted for Charlson comorbidity index, surgical procedure during hospital stay, duration of diabetes, and presence of malnutrition. D: Dynapenia; AO: Abdominal obesity; DAO: Dynapenic abdominal obesity; LOS: length of hospital stay; OR: odds ratio; CI: confidence interval; HR: hazard ratio. \*Logistic regression; † Cox regression.



**FIGURE 2.** Kaplan-Meier curve of 1-year mortality after hospital discharge, stratified by presence of DAO, in older patients with type 2 diabetes.

### DISCUSSION

In this prospective cohort study of 309 older patients with T2D, DAO was significantly associated with 2.5-fold odds of LOS ≥ 14 days and a 4.70-fold higher risk of 1-year mortality after hospital discharge (*vs.* patients without DAO).

The prevalence of dynapenia, AO, and DAO among our participants with T2D was 27.5, 30.7, and 32.4%,

respectively. Overall, 50.5% of the sample was female; of these women, 62% were diagnosed with DAO. A cross-sectional study of 20,198 older adults reported similar findings: 54% of that sample comprised women and the prevalence of DAO among them was 74.3%.<sup>22</sup> When we evaluated the main lifestyle factors of our patients, most were overweight (~70%) and sedentary (~85%). It has recently been shown that highly sedentary behavior may increase the risk of DAO in older adults.<sup>22</sup> In fact, sedentariness could potentially reduce muscle strength due to a lack of muscular contractility and other physiological consequences on muscle metabolism (e.g., reduced muscle glucose).<sup>23</sup> This is very useful information for individuals with T2D. However, in our participants there was no association between sedentary lifestyle and DAO. This can be explained by the fact that, despite their sedentarism, patients had reasonably on-target HbA1C levels (median of 7%). Additionally, those with a BMI ≥ 25 kg/m<sup>2</sup> had a higher prevalence of AO (44.7%) and DAO (30%). On the other hand, individuals with dynapenia had a higher prevalence of malnutrition (24.5%). A study of older persons with T2D found a prevalence of dynapenia of approximately 14%, increasing with longer time since diagnosis of diabetes.<sup>10</sup> This study observed a median (IQR) duration of diabetes of 10 (2 – 15) years and significant associations of dynapenia and DAO with time since diagnosis of diabetes (*p* < 0.05).

One of the main findings of this study is the association of dynapenia and DAO with prolonged hospitalization. Older patients with T2D who also had dynapenia or DAO were at 1.61-fold and 2.5-fold higher risk of LOS ≥ 14 days, respectively. Evidence from persons without diabetes corroborates our findings.<sup>12,13</sup> A study carried out



with 600 hospitalized individuals found that those with dynapenia had a 1.2-fold increase in their odds of prolonged hospitalization.<sup>12</sup> In a cohort study of 1136 individuals, dynapenia was associated with a LOS of 13 days when compared to those who had no dynapenia, regardless of age, sex, BMI, type of surgery, frailty, and nutritional status.<sup>13</sup> Dynapenia can contribute significantly to LOS.<sup>24</sup> Low muscle strength can limit ability to mobilize early, and lack of mobility can prolong the time needed to regain functional independence and ability to perform the activities of daily living, which can delay hospital discharge.<sup>3,24,25</sup> Moreover, it can increase the risk of falls and injuries during rehabilitation, which can lead to additional complications and further prolong LOS.<sup>26</sup> Indeed, a recent large longitudinal study showed that the presence of dynapenia and DAO accelerated the progression of frailty in older people.<sup>27</sup> Furthermore, the AO component of DAO affects muscle growth and metabolism by increasing the body fat percentage, decreasing muscle strength and muscle mass.<sup>27</sup> In older adults, intramuscular fat accumulation is inversely associated with muscle function.<sup>27</sup>

Another important finding from this study refers to the association between DAO and 1-year mortality after discharge. Older patients with DAO showed a 4.51-fold risk of mortality compared to those without DAO. However, DAO showed no significant association with in-hospital mortality, which could be explained by the relatively low number of deaths during hospitalization (6.1%). The InCHIANTI study evaluated the predictive value of DAO for incident disability, hospitalization, and mortality in 370 older men and 476 older women.<sup>15</sup> After adjustment for potential confounders, the relative risk of death was 1.47 (95%CI 1.09 – 1.97) for dynapenia/DAO when compared to the no-dynapenia/DAO group.<sup>15</sup> The Longitudinal Study of Aging, which involved 7030 participants with an 8-year follow-up, evaluated whether DAO was associated with cardiovascular mortality.<sup>28</sup> Individuals with dynapenia/AO and DAO showed a significantly higher risk of cardiovascular mortality.<sup>28</sup> DAO has also been significantly associated with a higher risk of incident multimorbidity, including diabetes.<sup>29</sup>

This study explored the combined effect of muscle weakness (dynapenia) and AO on adverse clinical outcomes in older adults with T2D.

Some key implications of the results of this study include:

1. Increased risk of prolonged LOS: the association between DAO and prolonged hospital stays suggests that older adults with both conditions may face more severe health complications, leading to

longer recovery times and increased healthcare resource utilization;

2. Higher 1-year mortality: the finding that DAO is linked to higher 1-year mortality rates indicates a significant risk factor for premature death. This highlights a critical need for early identification and intervention in this population to improve survival outcomes; and
3. Comprehensive management approaches: our findings underscore the importance of comprehensive management strategies that address both muscle weakness and AO in older adults with T2D.

The analysis of older individuals with T2D and DAO and its association with prolonged hospitalization and mortality within 1 year after hospital discharge are the core strengths of this study. Other aspects may also be interpreted as such. First, the prospective cohort study design allowed us to observe outcomes over time, providing a clearer picture of the temporal relationship between DAO and negative outcomes. Second, we aimed to study a specific, vulnerable population (older adults with T2D), which could encourage further studies on personalized interventions and care strategies. Third, detailed data collection on muscle strength, AO, LOS, and mortality provides a robust dataset for analysis, increasing the reliability of the results. Fourth, the 1-year follow-up period allowed us to assess longer-term outcomes, providing valuable insights into the progression and impact of DAO over time.

Our study also had some limitations. As it was conducted at a single center, its findings may not be generalizable to other settings or populations with different demographic or clinical characteristics. Moreover, despite adjusting for various confounders, some unmeasured factors may have influenced the relationship between dynapenic abdominal obesity and the outcomes of interest. Finally, this study did not provide information on specific interventions that might mitigate the risks associated with DAO, limiting the applicability of its results to clinical practice.

## CONCLUSIONS

Older patients with T2D and DAO were more likely to have prolonged hospitalizations and 1-year mortality than those without DAO. However, further research is needed to understand the underlying mechanisms linking DAO with diabetes. The findings of this study stress the importance of addressing both muscle strength and OA in older people with T2D to improve their clinical outcomes.

## DECLARATIONS

### Acknowledgements

We would like to thank all participating patients, Financiamento e Incentivo à Pesquisa of the Hospital de Clínicas de Porto Alegre (Fipe/HCPA) for funding this study, and the National Council for Scientific and Technological Development (CNPq).

### Conflict of interest

The authors declare no conflicts of interest.

### Funding

This study was supported by Financiamento e Incentivo à Pesquisa do Hospital de Clínicas de Porto Alegre (grant number #150068 )

### Authors' contribution

Mileni Vanti Beretta: conceptualization, formal analysis, investigation, methodology, project administration, resources, writing – original draft. Ticiana da Costa Rodrigues: conceptualization, investigation, methodology, writing – review & editing. Thais Steemburgo: formal analysis, investigation, methodology, resources.

### Ethical approval and informed consent

The study protocol was approved by the relevant institutional Research Ethics Committee (approval #150068) and all participants provided written informed consent following the recommendations established by the Declaration of Helsinki.

### Data availability statement

The data that support the findings of this study are available from the corresponding author, TS, upon reasonable request.

### Reporting standards guidelines

The authors adopted STROBE (Strengthening the Reporting of Observational studies in Epidemiology, <https://www.strobe-statement.org/>).

## REFERENCES

- Dhawan D, Sharma S. Abdominal obesity, adipokines and non-communicable diseases. *J Steroid Biochem Mol Biol.* 2020;203:105737. <https://doi.org/10.1016/j.jsmb.2020.105737>
- Clark BC, Manini TM. What is dynapenia? *Nutrition.* 2012;28(5):495–503. <https://doi.org/10.1016/j.nut.2011.12.002>
- Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age Ageing.* 2011;40(4):423–9. <https://doi.org/10.1093/ageing/af1051>
- Máximo RO, Oliveira DC, Ramírez PC, Luiz MM, Souza AF, Delinocente MLB, et al. Dynapenia, abdominal obesity or both: which accelerates the gait speed decline most? *Age Ageing.* 2021;50(5):1616–25. <https://doi.org/10.1093/ageing/afab093>
- Lv D, Shen S, Chen X. Association between dynapenic abdominal obesity and fall risk in older adults. *Clin Interv Aging.* 2022;17:439–45. <https://doi.org/10.2147/CIA.S347053>
- Oba K, Tamura Y, Ishikawa J, Suzuki H, Fujiwara Y, Tachibana A, et al. Dynapenic abdominal obesity is associated with mild cognitive impairment in patients with cardiometabolic disease: a cross-sectional study. *BMC Geriatr.* 2022;22(1):255. <https://doi.org/10.1186/s12877-022-02948-1>
- Rossi AP, Fantin F, Caliarì C, Zoico E, Mazzali G, Zanardo M, et al. Dynapenic abdominal obesity as predictor of mortality and disability worsening in older adults: a 10-year prospective study. *Clin Nutr.* 2016;35(1):199–204. <https://doi.org/10.1016/j.clnu.2015.02.005>
- da Silva Alexandre T, Scholes S, Ferreira Santos JL, de Oliveira Duarte YA, de Oliveira C. Dynapenic abdominal obesity increases mortality risk among English and Brazilian older adults: a 10-year follow-up of the ELSA and SABE studies. *J Nutr Health Aging.* 2018;22(1):138–44. <https://doi.org/10.1007/s12603-017-0966-4>
- Kalyani RR, Metter EJ, Egan J, Golden SH, Ferrucci L. Hyperglycemia predicts persistently lower muscle strength with aging. *Diabetes Care.* 2015;38(1):82–90. <https://doi.org/10.2337/dc14-1166>
- Mori H, Kuroda A, Ishizu M, Ohishi M, Takashi Y, Otsuka Y, et al. Association of accumulated advanced glycation end products with a high prevalence of sarcopenia and dynapenia in patients with type 2 diabetes. *J Diabetes Investig.* 2019;10(5):1332–40. <https://doi.org/10.1111/jdi.13014>
- Chang CY, Chu NF, Lin MH, Wang SC, Wu DM, Tsai MK, et al. Association between grip strength, obesity, and cardiometabolic risk factors among the community-dwelling elderly population in Taiwan. *Int J Environ Res Public Health.* 2022;19(18):11359. <https://doi.org/10.3390/ijerph191811359>
- Chites VS, Teixeira PP, Lima J, Burchell CF, Pereira TG, Silva FM. Reduced handgrip strength in hospital admission predicts prolonged hospital stay and death but is not accurate to identify malnutrition: a longitudinal study of reduced handgrip strength in hospitalized patients. *JPEN J Parenter Enteral Nutr.* 2021;45(5):1016–22. <https://doi.org/10.1002/jpen.1976>
- Marano L, Carbone L, Poto GE, Gambelli M, Noudem LLN, Grassi G, et al. Handgrip strength predicts length of hospital stay in an abdominal surgical setting: the role of frailty beyond age. *Aging Clin Exp Res.* 2022;34(4):811–7. <https://doi.org/10.1007/s40520-022-02121-z>
- McNicholl T, Curtis L, Dubin JA, Mourtzakis M, Nasser R, Laporte M, et al. Handgrip strength predicts length of stay and quality of life in and out of hospital. *Clin Nutr.* 2020;39(8):2501–9. <https://doi.org/10.1016/j.clnu.2019.11.006>
- Rossi AP, Bianchi L, Volpato S, Bandinelli S, Guralnik J, Zamboni M, et al. Dynapenic abdominal obesity as a predictor of worsening disability, hospitalization, and mortality in older adults: results from the InCHIANTI study. *J Gerontol A Biol Sci Med Sci.* 2017;72(8):1098–104. <https://doi.org/10.1093/gerona/glw203>
- Beretta MV, Dantas Filho FF, Freiberg RE, Feldman JV, Nery C, Rodrigues TC. Sarcopenia and type 2 diabetes mellitus as predictors of 2-year mortality after hospital discharge in a cohort of hospitalized older adults. *Diabetes Res Clin Pract.* 2020;159:107969. <https://doi.org/10.1016/j.diabres.2019.107969>
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987;40(5):373–83. [https://doi.org/10.1016/0021-9681\(87\)90171-8](https://doi.org/10.1016/0021-9681(87)90171-8)



18. World Health Organization. Physical status the use and interpretation of anthropometrics: report of a WHO expert committee. Geneva: WHO Technical Report Series; 1995.
19. Guigoz Y, Vellas B, Garry PJ. Assessing the nutritional status of the elderly: the mini nutritional assessment as part of the geriatric evaluation. *Nutr Rev.* 1996;54(1 Pt 2):S59-65. <https://doi.org/10.1111/j.1753-4887>
20. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing.* 2019;48(4):601. <https://doi.org/10.1093/ageing/afz046>
21. Brucki SMD, Nitrin R, Caramelli P, Bertolucci PHF, Okamoto IH. Suggestions for utilization of the mini-mental state examination in Brazil. *Arq Neuropsiquiatr.* 2003;61(3B):777-81. <https://doi.org/10.1590/s0004-282x2003000500014>
22. Smith L, López Sánchez GF, Rahmati M, Tully MA, Pizzol D, Veronese N, et al. Association between sedentary behavior and dynapenic abdominal obesity among older adults from low- and middle-income countries. *Aging Clin Exp Res.* 2024;36(1):109. <https://doi.org/10.1007/s40520-024-02763-1>
23. Keevil VL, Wijndaele K, Luben R, Sayer AA, Wareham NJ, Khaw KT. Television viewing, walking speed, and grip strength in a prospective cohort study. *Med Sci Sports Exerc.* 2015;47(4):735-42. <https://doi.org/10.1249/MSS.0000000000000453>
24. Vaishya R, Misra A, Vaish A, Ursino N, D'Ambrosi R. Hand grip strength as a proposed new vital sign of health: a narrative review of evidence. *J Health Popul Nutr.* 2024;43(1):7. <https://doi.org/10.1186/s41043-024-00500-y>
25. Kao CY, Su YC, Chang SF. The relationship between dynapenic abdominal obesity and fall: a systematic review and meta-analysis of 15,506 middle to older adults. *J Clin Med.* 2023;12(23):7253. <https://doi.org/10.3390/jcm12237253>
26. Akima H, Yoshiko A, Tomita A, Ando R, Saito A, Ogawa M, et al. Relationship between quadriceps echo intensity and functional and morphological characteristics in older men and women. *Arch Gerontol Geriatr.* 2017;70:105-11. <https://doi.org/10.1016/j.archger.2017.01.014>
27. Sun B, Wang J, Wang Y, Xiao W, Liu Y, Wang Y, et al. Associations of dynapenic abdominal obesity and frailty progression: evidence from two nationwide cohorts. *Nutrients.* 2024;16(4):518. <https://doi.org/10.3390/nu16040518>
28. Ramírez PC, Oliveira DC, Máximo RO, Souza AF, Luiz MM, Delinocente MLB, et al. Is dynapenic abdominal obesity a risk factor for cardiovascular mortality? A competing risk analysis. *Age Ageing.* 2023;52(1):afac301. <https://doi.org/10.1093/ageing/afac301>
29. Veronese N, Koyanagi A, Soysal P, Sapienza V, Ragusa FS, Bolzetta F, et al. Dynapenic abdominal obesity and incident multimorbidity: findings from the English longitudinal study on ageing. *Aging Clin Exp Res.* 2023;35(8):1671-8. <https://doi.org/10.1007/s40520-023-02455-2>