








New exercise programs for older adults in the community: evidence and applicability in the Latin American context

Novos programas de exercícios para idosos na comunidade: evidências e aplicabilidade no contexto latino-americano

Walter Sepúlveda-Loyola^a , Yshoner Antonio Silva-Díaz^b , Denilson de Castro Teixeira^c ,
Carol Saldías Solís^d , Danúbia da Cunha de Sá Caputo^e , Osvaldo Enrique Artaza Barrios^a ,
Alejandro Álvarez-Bustos^f 

^aFaculty of Health and Social Sciences,
Universidad de Las Américas — Santiago, Chile.

^bInstituto de Salud Integral Intercultural,
Facultad de Ciencias de la Salud, Universidad
Nacional Toribio Rodríguez de Mendoza —
Chachapoyas, Peru.

^cPrograma de Mestrado e Doutorado em
Ciências da Reabilitação, Universidade Estadual
de Londrina — Londrina (PR), Brazil.

^dEscuela de Kinesiología, Facultad de Salud,
Universidad Santo Tomás — Temuco, Chile.

^eLaboratório de Vibrações Mecânicas e Práticas
Integrativas, Departamento de Biofísica e
Biometria, Instituto de Biologia Roberto
Alcântara Gomes and Policlínica Piquet
Carneiro, Universidade do Estado do Rio de
Janeiro — Rio de Janeiro (RJ), Brazil.

^fCentro de Investigación Biomédica en Red
sobre Fragilidad y Envejecimiento Saludable,
Instituto de Salud Carlos III — Madrid, Spain.

Correspondence data

Walter Sepúlveda-Loyola — Echaurren, 140 —
Centro — 8370065 — Santiago, Chile.
E-mail: wsepulveda@udla.cl

Received on: March 23, 2025

Accepted on: March 23, 2025

How to cite this article: Sepúlveda-Loyola W, Silva-Díaz YA, Teixeira DC, Solís CS, Caputo DCS, Barrios OEA, et al. New exercise programs for older adults in the community: evidence and applicability in the Latin American context. *Geriatr Gerontol Aging*. 2025;19:e0000306.
https://doi.org/10.53886/gga.e0000306_EN

Copyright: © 2025 Sepúlveda-Loyola et al. This open-access article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Latin America is experiencing one of the fastest aging demographics globally. Projections indicate that more than 15% of its population will be over 60 years by 2030, and this figure will surpass 30% by 2060.¹ This demographic transition is accompanied by a significant rise in non-communicable diseases, such as diabetes, hypertension, and cardiovascular diseases.² These diseases have shown an exponential increase worldwide, particularly in Latin America. For example, the prevalence of diabetes in the region is expected to exceed 10% by 2050.²

As a consequence of these chronic conditions, sarcopenia and frailty have emerged as major health concerns among older adults, leading to functional decline and higher health care costs.^{3,4} Additionally, sedentary behavior, poor dietary habits, and the lack of structured physical activity programs have significantly contributed to the rising prevalence of comorbidities in older adults in Latin America.⁵ Given this rapid increase, it is imperative to develop and implement comprehensive strategies focused on preventive and therapeutic interventions to promote healthy aging in this region.

Promoting structured exercise programs tailored to the needs of older adults is one of the most effective strategies for mitigating the impact of chronic diseases and associated geriatric syndromes. Exercise is widely recognized for its role in improving cardiovascular health, glycemic control, muscle mass and function, functional independence, and overall quality of life.⁶ Exercise interventions for older populations encompass a wide range of modalities, including aerobic, resistance, coordination, balance, and flexibility exercises. Expert consensus guidelines recommend that these modalities be prescribed at varying intensities, adjusting for specific therapeutic objectives.⁶

For example, for individuals with reduced aerobic capacity and muscle strength, such as older adults living with cardiovascular or pulmonary diseases, aerobic and resistance training are the most commonly used exercise modalities.⁷ In contrast, resistance, balance, and gait exercises are prioritized for those at risk of falls or frailty.⁶ Therefore, the selection of exercise modalities and intensity should align with the therapeutic objective, considering the individual's functional status and existing comorbidities. Another critical consideration is the monitoring process during exercise sessions. While some older adults may require individualized supervision and tailored exercise programs, others may benefit from community-based exercise programs. Although group sessions provide less individualized attention and pose challenges in monitoring each participant, they offer significant

advantages, including fostering social connections, enhancing social interaction and engagement, and allowing for the inclusion of more participants per session.⁸

Cognitive function is also an essential factor to consider, since cognitive impairment may hinder the proper understanding or performance of tasks. Moreover, the ability to perform motor and cognitive tasks simultaneously declines with aging and is further impaired by certain comorbidities.⁹ Consequently, new exercise programs incorporating dual-task activities have been developed, including functional exercises designed to maintain both physical and cognitive health.¹⁰

Community-based exercise interventions offer a cost-effective solution for promoting physical activity among older adults, particularly in low-resource settings.¹¹ In Latin America, 89% of the community-based exercise programs feature traditional physical exercises, while 11% incorporate alternative programs such as Tai Chi and Pilates.¹² Several of these exercise programs are developed locally,^{8,12,13} while others are adapted from international models.^{14,15} One example is the VIVIFRAIL program,¹³ a multicomponent physical exercise initiative designed to prevent frailty and falls. It uses functional capacity assessments, such as the Short Physical Performance Battery (SPPB), to tailor exercise prescriptions for unsupervised sessions. In addition, a standardized functional exercise circuit (FEC) method has been successfully implemented in countries such as Chile and Brazil.^{8,12,13} These programs use structured multicomponent exercises, including aerobic capacity, strength, resistance, flexibility, coordination, and balance, designed for public spaces, parks, and community centers. Beyond structured exercise programs, lifestyle modifications play a crucial role in healthy aging.

Encouraging daily movement, reducing sedentary time, and promoting social activities that incorporate physical exertion can contribute to long-term well-being. In this regard, the VAMOS program (*Vida Ativa Melhorando a Saúde*) has yielded positive results in Brazil, demonstrating its effectiveness in community-based interventions for both middle-aged and older adults.^{14,15}

Despite the evidence published in this field, significant challenges remain unresolved. A recent systematic review found that most evidence on exercise programs in Latin America originates from only a few countries — primarily Brazil, Chile, Mexico, and Ecuador —, with most of them being implemented in Brazil.¹² In addition, exercise programs must be adaptable to different settings and individual conditions. Programs must be designed to be feasible, scalable, and accessible to ensure their effectiveness in diverse socioeconomic and geographic contexts, considering the peculiarities of rural communities or low-resource settings across Latin America.¹¹ For this reason, in this Special Issue, we invite researchers and clinicians from across Latin America to share their experiences with innovative exercise interventions aimed at promoting healthy aging and preventing comorbidities, geriatric syndromes, and their consequences. We encourage submissions that explore evidence-based strategies, including exercise programs in different settings (community-based, nursing homes, inpatient care, etc.), digital health innovations, exercise prescription and customization, and barriers and facilitators of exercise participation among older adults in Latin America. Through collaboration and knowledge exchange, we aim to advance scalable and sustainable interventions that promote active aging and enhance older adults' quality of life across this region.

DECLARATIONS

Authors' contribution

Walter Sepúlveda-Loyola: conceptualization, writing — original draft, writing — review & editing. Yshoner Antonio Silva-Díaz: writing — original draft, writing — review & editing. Denilson de Castro Teixeira: writing — review & editing. Carol Saldías Solís: conceptualization, writing — review & editing. Danúbia da Cunha de Sá Caputo: project administration, writing — review & editing. Osvaldo Enrique Artaza Barrios: conceptualization, writing — original draft. Alejandro Álvarez-Bustos: visualization, writing — review & editing

REFERENCES

1. Economic Commission for Latin America and the Caribbean. Ageing in Latin America and the Caribbean: inclusion and rights of older persons [Internet]. Santiago: ECLAC; 2022. Available from: <https://www.cepal.org/en/publications/48568-ageing-latin-america-and-caribbean-inclusion-and-rights-older-persons>. Accessed in Feb. 27, 2025.
2. Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, et al. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD 2019 study. *J Am Coll Cardiol*. 2020;76(25):2982–3021. <https://doi.org/10.1016/j.jacc.2020.11.010>

3. Álvarez-Bustos A, Rodríguez-Sánchez B, Carnicero-Carreño JA, Sepúlveda-Loyola W, García-García FJ, Rodríguez-Mañas L. Healthcare cost expenditures associated to frailty and sarcopenia. *BMC Geriatr.* 2022;22(1):747. <https://doi.org/10.1186/s12877-022-03439-z>
4. Da Mata FAF, Pereira PPDS, Andrade KRC, Figueiredo ACMG, Silva MT, Pereira MG. Prevalence of frailty in Latin America and the Caribbean: a systematic review and meta-analysis. *PLoS One.* 2016;11(8):e0160019. <https://doi.org/10.1371/journal.pone.0160019>
5. Strain T, Flaxman S, Guthold R, Semanova E, Cowan M, Riley LM, et al. National, regional, and global trends in insufficient physical activity among adults from 2000 to 2022: a pooled analysis of 507 population-based surveys with 5·7 million participants. *Lancet Glob Health.* 2024;12(8):e1232-43. [https://doi.org/10.1016/S2214-109X\(24\)00150-5](https://doi.org/10.1016/S2214-109X(24)00150-5)
6. Izquierdo M, Merchant RA, Morley JE, Anker SD, Aprahamian I, Arai H, et al. International Exercise Recommendations in Older Adults (ICFSR): expert consensus guidelines. *J Nutr Health Aging.* 2021;25(7):824-53. <https://doi.org/10.1007/s12603-021-1665-8>
7. Venkatesan P. GOLD COPD report: 2024 update. *Lancet Respir Med.* 2024;12(1):15-6. [https://doi.org/10.1016/S2213-2600\(23\)00461-7](https://doi.org/10.1016/S2213-2600(23)00461-7)
8. Loyola WS, Camillo CA, Torres CV, Probst VS. Effects of an exercise model based on functional circuits in an older population with different levels of social participation. *Geriatr Gerontol Int.* 2018;18(2):216-23. <https://doi.org/10.1111/ggi.13167>
9. Pasten JG, Aguayo JC, Aburto J, Araya-Quintanilla F, Álvarez-Bustos A, Valenzuela-Fuenzalida JJ, et al. Dual-task performance in individuals with chronic obstructive pulmonary disease: a systematic review with meta-analysis. *Pulm Med.* 2024;2024:1230287. <https://doi.org/10.1155/2024/1230287>
10. Varela-Vásquez LA, Minobes-Molina E, Jerez-Roig J. Dual-task exercises in older adults: a structured review of current literature. *J Frailty Sarcopenia Falls.* 2020;5(2):31-7. <https://doi.org/10.22540/JFSF-05-031>
11. Ibanez A, Maito M, Botero-Rodríguez F, Fittipaldi S, Coronel C, Migeot J, et al. Healthy aging meta-analyses and scoping review of risk factors across Latin America reveal large heterogeneity and weak predictive models. *Nat Aging.* 2024;4(8):1153-65. <https://doi.org/10.1038/s43587-024-00648-6>
12. Vásquez-Araneda E, Solís-Vivanco RI, Mahecha-Matsudo S, Zapata-Lamana R, Cigarroa I. Characteristics of physical exercise programs for older adults in Latin America: a systematic review of randomized controlled trials. *Int J Environ Res Public Health.* 2021;18(6):2812. <https://doi.org/10.3390/ijerph18062812>
13. Sepúlveda-Loyola W, Tricánico Maciel RP, Castro Teixeira D, Araya-Quintanilla F, Silva Junior RA, Álvarez-Bustos A, et al. Circuito de ejercicio funcional con tareas duales sobre variables clínicas relacionados con la sarcopenia. *Retos.* 2025;63:459-71. <https://doi.org/10.47197/retos.v63.110528>
14. Izquierdo M. Prescripción de ejercicio físico. El programa Vivifrail como modelo. *Nutr Hosp.* 2019;36(2):50-6. <https://doi.org/10.20960/nh.02680>
15. Benedetti T, Manta S, Gomez L, Rech C. Logical model of a behavior change program for community intervention — Active Life Improving Health — VAMOS. *Rev Bras Ativ Fis Saúde.* 2017;22(3):309-13. <https://doi.org/10.12820/rbafs.v.22n3p309-313>